



240 15
30 0, 25, 17, 18 . D
..... A ≥ 5
..... 18 3
..... 19, 20
..... 12.0
..... 2, 5

8. World Health Organization. Early childhood fact sheet. 2009. <http://www.who.int/mediacentre/factsheets/fs311/en/>. Accessed 15 September 2017.
9. World Health Organization. Report of the commission on ending childhood obesity. 2016. http://apps.who.int/iris/bitstream/10665/204176/1/9789241510066_eng.pdf. Accessed 15 September 2017.
10. Tremblay MS, Connor GS. Canadian health measures survey: brief overview. *Can J Public Health*. 2007;98:453–6.
11. Statistics Canada. Instructions for combining multiple cycles of Canadian health measures survey (CHMS) data. December 2015. http://www23.statcan.gc.ca/imdb-bmdi/document/5071_D4_T9_V2-eng.htm. Accessed 15 September 2017.
12. Day B, Langlois R, Tremblay M, Knoppers BM. Canadian health measures survey: ethical, legal and social issues. *Health Rep*. 2007;18:37–51.
13. Statistica Canada. Canadian health measures survey (CHMS) data user guide: cycle 2. Ottawa: Statistics Canada; 2013.
14. Statistics Canada. Canadian health measures survey (CHMS) data user guide: cycle 3. Ottawa: Statistics Canada; 2015.
15. Statistics Canada. Canadian health measures survey (CHMS) data user guide: cycle 4. Ottawa: Statistics Canada; 2017.
16. Statistics Canada. Canadian health measures survey. 2016. www.statcan.gc.ca/chms. Accessed 15 September 2017.
17. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008;40:181–8.
18. Pfeiffer KA, Dowda M, McIver KL, Pate RR. Factors related to objectively measured physical activity in preschool children. *Pediatr Exerc Sci*. 2009;21:196–208.
19. Rich C, Geraci M, Griffiths L, Sera F, Dezateux C, Cortina-Borja M. Quality control methods in accelerometer data processing: defining minimum wear time. *PLoS One*. 2013;8:e67206.
20. Kang M, Bjornson K, Barreira TV, Ragan BG, Song K. The minimum number of days required to establish reliable physical activity estimates in children aged 2–15 years. *Physiol Meas*. 2014;35:2229–37.
21. Adolph AL, Puyau MR, Vohra FA, Nicklas TA, Zakeri IF, Butte NF. Validation of uniaxial and triaxial accelerometers for the assessment of physical activity in preschool children. *J Phys Act Health*. 2012;9:944–53.
22. Colley RC, Harvey A, Grattan KP, Adamo KB. Impact of accelerometer epoch length on physical activity and sedentary behaviour outcomes for preschool-aged children. *Health Rep*. 2014;25:3–9.
23. Colley RC, Garriguet D, Adamo KB, Carson V, Janssen I, Timmons BW, Tremblay MS. Physical activity and sedentary behavior during the early years in Canada: a cross-sectional study. *Int J Behav Nutr Phys Act*. 2013;10:54.
24. WHO Multicentre Growth Reference Study Group. WHO child growth standards based on length/height, weight and age. *Acta Paediatr Suppl*. 2006;450:76–85.
25. Canadian Society for Exercise Physiology. Canadian physical activity, fitness and lifestyle approach (CPAFLA). 3rd ed. Ottawa: The Society; 2003.
26. Rust KF, Rao JNK. Variance estimation for complex surveys using replication techniques. *Stat Methods Med Res*. 1996;5:283–310.
27. Cliff DP, McNeill J, Vella SA, Howard S, Santos R, Batterham M, et al. Adherence to 24-Hour Movement Guidelines for the Early Years and associations with social-cognitive development among Australian preschool children. *BMC Public Health*. 2017;17(5). [in press].